FACTS for FAMILIES

No. 21 July 2004

Psychiatric Medication for Children and Adolescents Part 1- How Medications Are Used

Medication can be an effective part of the treatment for several psychiatric disorders of childhood and adolescence. A doctor's recommendation to use medication often raises many concerns and questions in both the parents and the youngster. The physician who recommends medication should be experienced in treating psychiatric illnesses in children and adolescents. He or she should fully explain the reasons for medication use, what benefits the medication should provide, as well as possible risks and side effects and other treatment alternatives.

Psychiatric medication should not be used alone. The use of medication should be based on a comprehensive psychiatric evaluation and be one part of a comprehensive treatment plan.

Before recommending any medication, the child and adolescent psychiatrist interviews the youngster and makes a thorough diagnostic evaluation. In some cases, the evaluation may include a physical exam, psychological testing, laboratory tests, other medical tests such as an electrocardiogram (EKG) or electroencephalogram (EEG) , and consultation with other medical specialists.

Medications which have beneficial effects may also have side effects, ranging from just annoying to very serious. As each youngster is different and may have individual reactions to medication, close contact with the treating physician is recommended. Do not stop or change a medication without speaking to the doctor. Psychiatric medication should be used as part of a comprehensive plan of treatment, with ongoing medical assessment and, in most cases, individual and/or family psychotherapy. When prescribed appropriately by a psychiatrist (preferably a child and adolescent psychiatrist), and taken as prescribed, medication may reduce or eliminate troubling symptoms and improve the daily functioning of children and adolescents with psychiatric disorders.

Medication may be prescribed for psychiatric symptoms and disorders, including, but not limited to:

- 1. **Bedwetting**-if it persists regularly after age 5 and causes serious problems in low self-esteem and social interaction.
- 2. **Anxiety** (school refusal, phobias, separation or social fears, generalized anxiety, or posttraumatic stress disorders)-if it keeps the youngster from normal daily activities.

- 3. **Attention deficit hyperactivity disorder (ADHD)** -marked by a short attention span, trouble concentrating and restlessness. The child is easily upset and frustrated, often has problems getting along with family and friends, and usually has trouble in school.
- 4. **Obsessive-compulsive disorder (OCD)** -recurring obsessions (troublesome and intrusive thoughts) and/or compulsions (repetitive behaviors or rituals such as handwashing, counting, checking to see if doors are locked) which are often seen as senseless but which interfere with a youngster's daily functioning.
- Depression-lasting feelings of sadness, helplessness, hopelessness, unworthiness
 and guilt, inability to feel pleasure, a decline in school work and changes in
 sleeping and eating habits.
- 6. **Eating disorder**-either self-starvation (anorexia nervosa) or binge eating and vomiting (bulimia), or a combination of the two.
- 7. **Bipolar (manic-depressive) disorder**-periods of depression alternating with manic periods, which may include irritability, "high" or happy mood, excessive energy, behavior problems, staying up late at night, and grand plans.
- 8. **Psychosis**-symptoms include irrational beliefs, paranoia, hallucinations (seeing things or hearing sounds that don't exist) social withdrawal, clinging, strange behavior, extreme stubbornness, persistent rituals, and deterioration of personal habits. May be seen in developmental disorders, severe depression, schizoaffective disorder, schizophrenia, and some forms of substance abuse.
- 9. **Autism**-(or other pervasive developmental disorder such as Asperger's Syndrome)-characterized by severe deficits in social interactions, language, and/or thinking or ability to learn, and usually diagnosed in early childhood.
- 10. **Severe aggression**-which may include assaultiveness, excessive property damage, or prolonged self-abuse, such as head-banging or cutting.
- 11. **Sleep problems**-symptoms can include insomnia, night terrors, sleep walking, fear of separation, anxiety.

For additional information about psychiatric medications see Facts for Families: #29 Psychiatric Medication for Children and Adolescents: Part II-Types of Medications #51 Psychiatric Medications for Children and Adolescents: Part III-Questions to Ask.

For additional information see Facts for Families: #00 Definition of a Child and Adolescent Psychiatrist #25 Know Where to Seek Help for Your Child #52 Comprehensive Psychiatric Evaluation

If you find *Facts for Families*[©] helpful and would like to make good mental health a reality, consider donating to the <u>Campaign for America's Kids</u>. Your support will help us continue to produce and distribute Facts for Families, as well as other vital mental health information, free of charge.

You may also mail in your contribution. Please make checks payable to the AACAP and send to *Campaign for America's Kids*, P.O. Box 96106, Washington, DC 20090.

The American Academy of Child and Adolescent Psychiatry (AACAP) represents over 8,500 child and adolescent psychiatrists who are physicians with at least five years of additional training beyond medical school in general (adult) and child and adolescent psychiatry.

PSYCHIATRIC MEDICATION FOR CHILDREN AND ADOLESCENTS-PART I, "Facts for Families,"

No. 21 (7/04)

Facts for Families© information sheets are developed, owned and distributed by AACAP. Hard copies of Facts sheets may be reproduced for personal or educational use without written permission, but cannot be included in material presented for sale or profit. All Facts can be viewed and printed from the AACAP website (www.aacap.org). Facts sheets may not be reproduced, duplicated or posted on any other website without written consent from AACAP. Organizations are permitted to create links to AACAP's website and specific Facts sheets. For all questions please contact the AACAP Communications & Marketing Coordinator, ext. 154.

If you need immediate assistance, please dial 911.

Copyright © 2012 by the American Academy of Child and Adolescent Psychiatry.